

Kayla Spicer and Joseph Goyette, Dr. David Thoms  
College of Arts and Sciences

## Abstract

Understanding how plants differentiate between harmful and beneficial bacteria is essential for advancing agricultural sustainability and plant health. Gaining faster insights into these defense mechanisms enables early disease detection and timely intervention, ultimately improving crop resilience. Additionally, harnessing beneficial microbes can enhance plant stress tolerance and support sustainable farming practices. The primary research question guiding this study is: "How do plants distinguish between harmful and beneficial microbes?" We hypothesize that membrane-associated proteins play a crucial role in this differentiation, particularly in detecting bacterial pore-forming toxins. To test this hypothesis, we utilized a microplate method where *Arabidopsis thaliana* seedlings were planted into 96-well plates. The roots were exposed to bacterial co-cultures, and plant-bacterial interactions were monitored over seven days. Bacterial growth was quantified using fluorescence plate readings. Our findings indicate that differences in microbial load exist across genotypes, particularly in mutants deficient in immune signaling components. These results provide insights into how plants recognize microbial threats and could inform future strategies for enhancing crop resistance.

## Introduction

Plants interact with both beneficial and pathogenic microbes in the rhizosphere, but how they distinguish between them, especially those producing pore-forming toxins, is poorly understood.

Research suggests membrane-associated proteins play a role in immune signaling, particularly in recognizing bacterial virulence factors.

This study investigates the role of synaptotagmin proteins in microbial recognition by analyzing how various *Arabidopsis* mutants respond to co-cultures of beneficial and pathogenic *Pseudomonas fluorescens* strains.

## Methods

*Arabidopsis thaliana* seedlings were grown in a 96-well microplate system designed by Melissa Chen, with roots suspended in microbial solutions while shoots remained above the surface.

The experiment included six plant genotypes:

- Controls:** *Col-0* (wild-type), *bak1/bkk1/cerk1* triple mutant
- Experimental mutants:** *syt1*, *syt5*, *NDR1*, and *SOBIR1* mutants

After five days of initial growth, plants were inoculated with bacterial co-cultures.


Plants were co-inoculated with two *Pseudomonas fluorescens* strains:

- WSC365* – a beneficial strain labeled with mNeonGreen fluorescence protein
- N2C3* – a pathogenic strain labeled with mScarlet fluorescence protein


Seedlings remained in bacterial culture for seven days.

Bacterial colonization was measured using a fluorescence plate reader. Fluorescence intensity was converted to bacterial concentration using a standard curve, allowing for a quantitative assessment of microbial load across plant genotypes.


Bacterial growth was analyzed comparatively across plant genotypes to assess the influence of immune signaling mutations on microbial interactions.



The concentrations of WSC365 (left) and N2C3 (right) are analyzed and compared across different mutants.



The overall bacterial concentration (left) and the ratio of N2C3 to WSC365 are compared across different mutants.



*Arabidopsis thaliana* at the time of inoculation (left) and seven days afterward (right).

## Results

Because plants were exposed to co-cultures of bacteria rather than isolated pathogens, differences in microbial colonization were influenced by immune-related mutations rather than a direct pathogen-only response.

Significant differences in bacterial load were observed between *syt1* and *syt5* mutants and other plant genotypes.

While overall bacterial presence remained similar across samples, notable variations in beneficial strain colonization were observed between genotypes.

## Discussion

Our findings suggest that *SYT5*, a synaptotagmin protein involved in vesicle trafficking, plays a crucial role in differentiating beneficial and harmful microbes.

The presence of pore-forming toxins (*PFTs*) in *N2C3* may contribute to these findings, as synaptotagmins potentially mediate membrane repair following toxin exposure.

*bak1/bkk1/cerk1* triple mutants served as immune signaling-deficient controls and exhibited expected colonization trends, reinforcing their role as a baseline comparison.

Future research should investigate the mechanistic relationship between synaptotagmins and microbial recognition using additional molecular assays.

## Resources

Cheng, Y. T., et al. (2024, September 6). Roles of microbiota in autoimmunity in *Arabidopsis* leaves. *Nature News*. Nature Publishing Group. <https://www.nature.com/articles/s41477-024-01779-9>

Li, J., et al. (2024, March 31). Association mechanism between *Arabidopsis* immune coreceptor BAK1 and *Pseudomonas syringae* effector HopF2. *Biochemical and Biophysical Research Communications*. Academic Press.

Lundberg, D. S., et al. (2012, August 1). Defining the core *Arabidopsis thaliana* root microbiome. *Nature News*. Nature Publishing Group.

Chen, M. Y., et al. (2024, April 29). Order among chaos: High throughput MYCroplanters can distinguish interacting drivers of host infection in a highly stochastic system. *bioRxiv*. Cold Spring Harbor Laboratory.

*Arabidopsis* Biological Resource Center. (n.d.). Seed stock information. The Ohio State University. <https://abrc.osu.edu>

## Acknowledgments

We would like to sincerely thank our research mentor, Dr. David Thoms, for his guidance and support, and The Center for Undergraduate Research and Engagement (CRE) and Undergraduate Research Opportunity Program (UROP) for providing us with this incredible research opportunity. Their support has been invaluable in our academic and research growth.