

Savannah Boutwell, Jessica Cassette, Allison Bowerfind, Jens Cuba, Isabela Fernandez, Allyson May, Katie Kirk, Justine Moss, Joseph Watso

INTRODUCTION

- Breathing patterns can influence the perception of physical effort and breathlessness at rest and during exercise (1). The rating of perceived exertion (RPE) may be lower during oral-only breathing than oronasal (i.e., mouth and nose) breathing during shuttle run test in young male adults (2).
- Nasal versus oral breathing reduces cardiorespiratory demands during exercise (3). However, it is unknown whether nasal versus oral breathing reduces the perception of physical effort and breathlessness in young male and female adults.

PURPOSE

- Therefore, the purpose of this study was to test the hypothesis that RPE and the rating of perceived breathlessness (RPB) would be lower with nasal versus oral breathing.

EXPERIMENTAL DESIGN

Part 1: Recruitment

- We enrolled 12 participants aged 18 - 19 years old with body mass index values of 17 - 26 kg/m².
- After receiving consent from the participant, we measured their body mass and height.

Part 2: Resting Measurements

- We randomized the order of nose-only and mouth-only breathing periods (5 minutes each) between participants. We used an auditory metronome to keep participants' respiratory rate consistent, based on their 'free breathing' respiratory rate, using a fabric respiration belt.
- Resting on a semi-recumbent bed, we attached equipment to monitor blood oxygen saturation, heart rate, and blood pressure (not reported on this poster).
- We asked participants to report a RPE (Figure 1) and RPB (Figure 2) at the end of the rest periods.

Part 3: Submaximal Exercise

- We used the same condition order as used during rest periods for two 7-minute bouts of exercise at 75 watts on a semi-recumbent cycle ergometer, with RPE and RPB reported in the last minute.

Rating of Perceived Exertion

6	No exertion
7	Extremely light
8	
9	Very light
10	
11	Light
12	
13	Somewhat hard
14	
15	Hard
16	
17	Very hard
18	
19	Extremely hard
20	Maximal exertion

Rating of Perceived Breathlessness

0	Nothing at all
0.5	Very, very light (just noticeable)
1	Very slight
2	Slight
3	Moderate
4	Somewhat severe
5	Severe
6	Very severe
7	
8	
9	Very, very severe (almost maximal)
10	Maximal

Figure 1:

Figure 2:

RESULTS




Figure 3: We found that resting RPB was higher during oral versus nasal breathing (B). However, we did not observe differences in the remaining variables at rest (A) or during exercise (C-D). Importantly, respiratory rate not different between conditions at rest (nasal: 17 ± 4 vs. oral: 16 ± 4 bpm, p=0.18) or during exercise (nasal: 26 ± 5 vs. oral: 24 ± 4 bpm, p=0.19). We used paired, two-tailed t-tests for exercise RPE and resting/exercise respiratory rate, and Wilcoxon matched-pairs signed rank tests for all other variables because they failed (p>0.05) the Shapiro-Wilk normality test.

CONCLUSIONS

These preliminary findings suggest that while nasal breathing at rest can reduce subjective ratings of breathlessness, it does not affect subjective ratings of exertion or breathlessness during submaximal exercise.

REFERENCES

- Dallam, G. ., and B. . Kies. "The Effect of Nasal Breathing Versus Oral and Oronasal Breathing During Exercise: A Review". *Journal of Sports Research*, vol. 7, no. 1, Jan. 2020, pp. 1-10, doi:10.18488/journal.90.2020.71.1.10.
- Meir, R., et al. (2014). "The acute effect of mouth only breathing on time to completion, heart rate, rate of perceived exertion, blood lactate, and ventilatory measures during a high-intensity shuttle run sequence." *J Strength Cond Res* 28(4): 950-957.
- Hall, R. L. (2005). "Energetics of nose and mouth breathing, body size, body composition, and nose volume in young adult males and females." *Am J Hum Biol* 17(3): 321-330.

ACKNOWLEDGMENTS

Supported in part by Florida State University startup funds (JCW), NIH K01HL160772 (JCW), and a UROP Materials Grant.