



Characterization of Deep-Sea Habitat Distribution on Two Seamounts of the Papahānaumokuākea Marine National Monument

Nicole Hernandez, Emily Irwin, Virginia Biede, Dr. Mauricio Silva, & Dr. Amy Baco-Taylor



Department of Earth Ocean and Atmospheric Sciences

Background

The North Pacific Ocean stretches from the Eastern Coast of Asia to the Western Coast of the North American continent. Undersea mountains formed from past volcanic eruptions, known as seamounts, make up nearly 28.8 square kilometers of Earth's surface (NOAA) with their highest abundance in the Pacific. Seamounts are most important for the biological richness they provide as underwater habitats. These seamounts feature different levels of rugosity, substrate types, and substrate sizes. These factors have a significant impact on seamount community composition, as do the dominant corals which also provide habitat to many different species (Rogers, A. 2018).

The Northwest Pacific Ocean has not been unaffected by global warming and human activity. It is also one of the most biologically diverse regions in the world hosting tens of thousands of marine species (Liao et al. 2018). The oceans are mostly made up of areas beyond national jurisdiction (ABNJ), which are key in biodiversity that is not possessed by States. Many ABNJ areas have deteriorated due to the lack of regulations on human activities (IUCN. 2023). Trawling in the Hawaiian Ridge and Emperor Seamount Chain in past years has led to near-total devastation of the benthic communities on parts of the underwater mountain range. Near entire ecosystems were depleted of corals and suffered as a consequence. Over years of protection, however, signs of life have begun to return despite their slow growth. It's been shown that over a 30-40 year period, measurable, but not total, recovery is possible.

Methods II



Pisces IV

Pisces V

Figure 1.1 and 2.2. Images of the battery-powered deep-submergence vehicles used to collect video footage of the seamounts underwater.

Statistics

Table with columns for Site, transect, C, CR, IC, DR, GS, UC, LR, M, RCA, RCA, S, WS. It contains statistical data for various sites and transects, with some cells highlighted in red to indicate significant differences.

Figure 2.1 - Shows the different p-values of each characteristic, highlighted text indicates significant differences.

Conclusion

After running a statistical two sample two tailed t-test which is used to compare rugosity levels, substrate composition, and substrate size on the Southeast Hancock and the Brook banks. We obtained a p-value (p>0.05) for all levels of rugosity, for most levels of substrate size, and for most substrate composition. With a p-value that is greater than our alpha=0.05 which means we reject the null hypothesis and there is a significant difference between substrate characteristics between the Southeast Hancock and Brooks Banks seamounts. Both seamounts have different characteristics that are able to sustain different ecosystems necessary for biodiversity. Brooks Banks has a lower level of rugosity which means not a lot of coral grows in this area since there's a low presence of boulders and larger substrates and mostly made up of sand where corals aren't able to grow. While Southeast Hancock has a larger presence of these bigger substrates which allows corals to thrive.

Results




Figure 3.1, 3.2, and 3.3 Overall percentage composition of rugosity, substrate size, and substrate on Southeast Hancock and Brooks Bank seamounts. Each column represents one XXXm long transect or Xxmin of video.




Figure 4.1. Rugosity levels on the Southeast Hancock seamount showed a trend of being higher compared to the Brooks Bank seamount with all levels having a p-value > 0.05, we can conclude that there is a significant difference between the characteristic of both seamounts.

Discussion


Biodiversity is essential for processes that support life on earth, without this one a large range of organisms would not be able to have ecosystems that provide essential sources to sustain human life. Our research showed the difference among seamounts that each one apport to the diversity of ecosystems. Corals are an essential system of our oceans, they provide homes, food, and overall support for the well-being of marine life. Corals are common in transects with higher depths and a higher presence of cobbles and boulders (Mortensen, P and Mortensen, L. 2004). South Hancock would be more likely to sustain a coral ecosystem due to the higher levels of rugosity. While other types of organisms thrive in habitats with low rugosity and smaller substrate composition. Where Brooks Bank has an advantage over these organisms. The Hawaiian Ridge and the emperor seamount ridge stretch across the North Pacific that support vast biodiversity. However, there is no legal mechanism that provides protection for some of these areas which are critical to the survival of these areas that are being harmed by human activity like trawling. The overall goal of our project as well as many other scientists is to determine the location of these patches to bring up to the United Nations to include as part of areas where human activity is not allowed to be able to protect such critical organisms.

Acknowledgments


This would have not been possible without the support of our mentors Virginia Biede, Dr. Amy Baco-Taylor, and Dr. Mauricio Silva, as well as the Pisces crew who collected all of the data by AUV and the crew of the Ka'imikai-O-Kanaloa. This research was supported by NSF Grant OCE-1334652 to ARB and OCE-1334675 to E. Brendan Roark and the FSU UROP program.

References

Colton, D. M., and M. Pinsk. 2021. Coral Reef Alliance. Local threats to coral reefs
Forney, K. A., E. Becker, D. G. Foley, J. Barlow, and E. M. Oleson. 2015. Habitat-based models of cetacean density and distribution in the central North Pacific. *Endangered Species Research* 27: 1-20.
Jiang, X., C. Dong, Y. Ji, C. Wang, Y. Shu, L. Liu, and J. Ji. 2021. Influences of Deep-Water Seamounts on the Hydrodynamic Environment in the Northwestern Pacific Ocean. *Journal of Geophysical Research: Oceans* 126 (12).
Madhav, K., S. Senaratna, S. Okayasu, and S. Wataru. 2018. Regional assessment report on biodiversity and ecosystem services for Asia and the Pacific. IPBES secretariat.
McQuaid, K. A., M. J. Attrill, M. R. Clark, A. Copley, A.G. Glover, C.R. Smith, and K.L. Howell. 2020. Using habitat classification to assess the representativity of a protected area network in a large, data-poor area targeted for deep-sea mining. *Frontiers*.
Therriault, T., C. Park, and J. Rice. 2016. *North Pacific Ocean*. United Nations.
2021. *Emperor seamounts*. NOAA.
Mortensen, P.B., and L. Buhl-Mortensen. 2004. Distribution of deep-water gorgonian corals in relation to benthic habitat features in the *Northeast Channel (Atlantic Canada)*. *Marine Biology* 144, 1223-1238.



Brooks Bank



Southeast Hancock

Methods I

Over the past decade, the destruction of deep-sea habitats has led to serious problems on earth. We wanted to look at two different seamounts found in the Hawaiian Ridge. Our study focused on Southeast Hancock and the Brooks Banks seamounts. The human-occupied vehicles named Pisces IV and Pisces V, were used to collect all the images. To analyze and collect data from these images we used two programs; Excel and ImageJ. In ImageJ, we uploaded the images and used the software to determine the area and rugosity of the substrate. Rugosity is the variations of amplitude in the height of a surface. We then proceeded to use a tool called "region of interest" (ROI) that would create fifteen randomized points to be categorized in terms of substrate composition and size. The rugosity scale ranged from 1-4 from low to high and the substrate size scale ranged from 1-8. This data was then collated in order to be statistically analyzed via a 2 tailed T-test.




Figure 5.1. Elevation maps of both Southeast Hancock and Brooks Banks, as well as the relative location of each seamount within the Papahānaumokuākea Marine National Monument.